

PARTITION REGULARITY WITHOUT THE COLUMNS PROPERTY

BEN BARBER, NEIL HINDMAN, IMRE LEADER, AND DONA STRAUSS

ABSTRACT. A finite or infinite matrix A with rational entries is called *partition regular* if whenever the natural numbers are finitely coloured there is a monochromatic vector x with $Ax = 0$. Many of the classical theorems of Ramsey Theory may naturally be interpreted as assertions that particular matrices are partition regular. In the finite case, Rado proved that a matrix is partition regular if and only it satisfies a computable condition known as the *columns property*. The first requirement of the columns property is that some set of columns sums to zero.

In the infinite case, much less is known. There are many examples of matrices with the columns property that are not partition regular, but until now all known examples of partition regular matrices did have the columns property. Our main aim in this paper is to show that, perhaps surprisingly, there are infinite partition regular matrices without the columns property — in fact, having no set of columns summing to zero.

We also make a conjecture that if a partition regular matrix (say with integer coefficients) has bounded row sums then it must have the columns property, and prove a first step towards this.

1. INTRODUCTION

Let A be an $u \times v$ matrix with rational entries. We say that A is *kernel partition regular*, or simply *partition regular*, if for every finite colouring of the natural numbers $\mathbb{N} = \{1, 2, \dots\}$ there is a monochromatic vector $\vec{x} \in \mathbb{N}^v$ with $A\vec{x} = \vec{0}$. In other words, A is partition regular if for every positive integer k , and every function $c : \mathbb{N} \rightarrow \{1, \dots, k\}$, there is a vector $\vec{x} = (x_1, \dots, x_v) \in \mathbb{N}^v$ with $c(x_1) = \dots = c(x_v)$ such that $A\vec{x} = \vec{0}$. We may also speak of the ‘system of equations $A\vec{x} = \vec{0}$ ’ being partition regular.

Many of the classical results of Ramsey Theory may naturally be considered as statements about partition regularity. For example, Schur’s Theorem [13], that in any finite colouring of the natural numbers we may solve $x + y = z$ in one colour class, is precisely the assertion that the 1×3 matrix $(1, 1, -1)$ is partition regular. As another example, the theorem of van der Waerden [15] that, for any m , every finite colouring of the natural numbers contains a monochromatic arithmetic progression with m terms, is (with the strengthening that we may also choose the common difference of the sequence to have the same colour) exactly the statement

2010 *Mathematics Subject Classification.* 05D10.

The second author acknowledges support received from the National Science Foundation (USA) via Grant DMS-1160566.

that the $(m-1) \times (m+1)$ matrix

$$\begin{pmatrix} 1 & 1 & -1 & 0 & \dots & 0 & 0 \\ 1 & 0 & 1 & -1 & \dots & 0 & 0 \\ 1 & 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 0 & 0 & 0 & \dots & 1 & -1 \end{pmatrix}$$

is partition regular.

In 1933 Rado characterized those matrices that are partition regular over \mathbb{N} in terms of the *columns property*.

Definition 1.1. Let $u, v \in \mathbb{N}$ and let A be a $u \times v$ matrix with entries from \mathbb{Q} . Denote the columns of A by $\langle \vec{c}_i \rangle_{i=0}^{v-1}$. The matrix A satisfies the *columns property* if and only if there exist $m \in \{1, 2, \dots, v\}$ and a partition $\langle I_t \rangle_{t=0}^{m-1}$ of $\{0, 1, \dots, v-1\}$ such that

- (1) $\sum_{i \in I_0} \vec{c}_i = \vec{0}$ and
- (2) for each $t \in \{1, 2, \dots, m-1\}$, $\sum_{i \in I_t} \vec{c}_i$ is a linear combination with coefficients from \mathbb{Q} of $\{\vec{c}_i : i \in \bigcup_{j=0}^{t-1} I_j\}$.

Theorem 1.2. Let $u, v \in \mathbb{N}$ and let A be a $u \times v$ matrix with entries from \mathbb{Q} . Then A is partition regular if and only if A satisfies the columns property.

Proof. [12, Satz IV]. □

In this paper, we are concerned with partition regularity of infinite systems of homogeneous linear equations or, equivalently, with partition regularity of infinite matrices. In all cases we shall assume that the equations we are dealing with have finitely many terms; that is, each row of the matrix of coefficients has all but finitely many entries equal to 0. The definition of partition regularity for a matrix (or a system of linear equations) is verbatim the same as for finite matrices or systems of equations.

The first example of a (non-trivial) infinite partition regular system of equations was constructed in 1974 [4], proving a conjecture of Graham and Rothschild: in any finite colouring of the natural numbers there is a sequence x_1, x_2, \dots of natural numbers such that the set

$$FS(x_1, x_2, \dots) = \left\{ \sum_{i \in I} x_i : I \subseteq \mathbb{N}, I \text{ finite and non-empty} \right\}$$

is monochromatic. This is also known as the Finite Sums Theorem. (It is worth remarking that the finite analogue of this, known as Folkman's Theorem, stating that, for any m , in any finite colouring of the natural numbers we may find x_1, \dots, x_m with $FS(x_1, \dots, x_m)$ monochromatic, follows easily from Rado's Theorem.) Since then, several other infinite partition systems have been found — for example, the Milliken–Taylor Theorem [11] [14] and several systems in [5]. See [6] for general background on this.

We take $\omega = \mathbb{N} \cup \{0\}$. The columns property has an obvious extension to infinite matrices.

Definition 1.3. Let $u, v \in \mathbb{N} \cup \{\omega\}$ and let A be a $u \times v$ matrix with entries from \mathbb{Q} . Denote the columns of A by $\langle \vec{c}_i \rangle_{i < v}$. The matrix A satisfies the *columns property* if and only if there exists a partition $\langle I_\sigma \rangle_{\sigma < \mu}$ of v , where $\mu \in \mathbb{N} \cup \{\omega\}$, such that

- (1) $\sum_{i \in I_0} \vec{c}_i = \vec{0}$ and
(2) for each $t \in \mu \setminus \{0\}$, $\sum_{i \in I_t} \vec{c}_i$ is a linear combination with coefficients from \mathbb{Q} of $\{\vec{c}_i : i \in \bigcup_{j < t} I_j\}$.

We stress that the sums in this definition are *not* required to be finite. Note that infinite sums of columns do always make sense, because our matrices have only finitely many non-zero entries in each row. It is also worth remarking that if one insisted on finite sums then nothing would work: even the matrix corresponding to the Finite Sums Theorem needs infinite sums for the columns property.

It is easy to see that, for infinite matrices, the columns property is not sufficient for partition regularity. Consider for example the system of equations $x_n - x_{n+1} = y_n$, $n \in \omega$. As a matrix equation this corresponds to

$$\begin{pmatrix} 1 & -1 & -1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 1 & -1 & -1 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 1 & -1 & -1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \\ x_1 \\ y_1 \\ \vdots \end{pmatrix} = \vec{0}.$$

The system of equations is not partition regular over \mathbb{N} for the trivial reason that it has no solutions there at all; any solution must have $x_n > x_{n+1}$ for each n . To see that the matrix satisfies the columns property, let $I_0 = \{0, 2, 4, 6, \dots\}$ and $I_1 = \{1, 3, 5, \dots\}$. Then $\sum_{i \in I_0} \vec{c}_i = \vec{0}$ and $\sum_{i \in I_1} \vec{c}_i = \sum_{t=1}^{\infty} t\vec{c}_{2t}$.

We shall show in Section 2 of this paper there is a system of linear equations which is partition regular over \mathbb{N} but for which the coefficient matrix has no set of columns summing to zero. We mention in passing that the entries and the columns of this matrix will be ‘nicely behaved’: all entries are from the set $\{-1, 0, 1, 2\}$ and each column of the coefficient matrix has at most three nonzero entries.

The example mentioned in the previous paragraph has the property that the row sums of the coefficient matrix are unbounded. We also show in Section 2 that if A is a matrix which is partition regular over \mathbb{N} and the sums of the absolute values of the entries in any row are bounded, then there must be some set of columns that sum to zero. We do not know whether such a matrix must satisfy the full columns property.

The proof that the system mentioned above is partition regular over \mathbb{N} depends on the following lemma, which is based on results in [2]. The lemma refers to *central* sets. From the point of view of the partition regularity of our system of equations, central sets have two key properties.

- Whenever \mathbb{N} is finitely coloured, one of the colour classes must be central.
- Any finite partition regular system of homogeneous linear equations has solutions in any central set (see [10, Theorem 15.16(b)]). (The corresponding statement is not true for infinite systems.)

In the statement of this lemma, $m \cdot \mathbb{Z} = \{m \cdot n : n \in \mathbb{Z}\}$ while $kC = C + C + \dots + C$ (k times).

Lemma 1.4. *Let C be a central subset of \mathbb{N} . There exist m and K in \mathbb{N} such that if $k \geq K$, then $m \cdot \mathbb{Z} \subseteq C - kC$.*

Section 3 consists of a proof of (a generalisation of) Lemma 1.4, and also a strengthening of the main result of Section 2 (namely Theorem 2.1). The proofs

depend on the algebraic structure of the Stone-Ćech compactification $\beta\mathbb{N}$ of the discrete space \mathbb{N} . We give a brief introduction to that structure in that section. We mention that the point of Lemma 1.4, and the point of using central sets, is that then the proof of Theorem 2.1 is clean and direct.

2. PARTITION REGULARITY DOES NOT IMPLY THE COLUMNS PROPERTY

In this section we consider the following system of equations:

$$2x_n + x_{2^n} + x_{2^{n+1}} + \cdots + x_{2^{n+1}-1} = y_n, \quad n \in \omega$$

The simplest way to represent the matrix of coefficients is as an $\omega \times (\omega + \omega)$ matrix. That is, we let A be the $\omega \times \omega$ matrix such that, for $i, j \in \omega$,

$$a_{i,j} = \begin{cases} 2 & \text{if } j = i, \\ 1 & \text{if } 2^i \leq j < 2^{i+1}, \\ 0 & \text{otherwise,} \end{cases}$$

so that

$$A = \begin{pmatrix} 2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \cdots \\ 0 & 2 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & \cdots \\ 0 & 0 & 2 & 0 & 1 & 1 & 1 & 1 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Letting I be the $\omega \times \omega$ identity matrix, we then have that the matrix of coefficients for our system of equations is $(A \quad -I)$. It is immediate that there is no non-empty set of columns of this matrix summing to $\vec{0}$.

Theorem 2.1. *For any central subset C of \mathbb{N} , there exist \vec{x} and \vec{y} in C^ω such that $(A \quad -I) \begin{pmatrix} \vec{x} \\ \vec{y} \end{pmatrix} = \vec{0}$. In particular the system of equations*

$$2x_n + x_{2^n} + x_{2^{n+1}} + \cdots + x_{2^{n+1}-1} = y_n, \quad n \in \omega$$

is partition regular over \mathbb{N} .

Proof. Pick m and K as guaranteed for C by Lemma 1.4. Pick $M \in \mathbb{N}$ such that $2^{M+1} \geq m - 2 + K$. Let P be the $(M+1) \times 2^{M+1}$ matrix consisting of rows $0, 1, \dots, M$ and columns $0, 1, \dots, 2^{M+1} - 1$ of A and let I_{M+1} be the $(M+1) \times (M+1)$ identity matrix. Then $(P \quad -I_{M+1})$ satisfies the columns property. Since, as we remarked earlier, any finite partition regular system of homogeneous linear equations has solutions in any central set, we may choose $\vec{x} = \langle x_i \rangle_{i=0}^{2^{M+1}-1}$ and $\vec{y} = \langle y_i \rangle_{i=0}^M$ in C such that $(P \quad -I_{M+1}) \begin{pmatrix} \vec{x} \\ \vec{y} \end{pmatrix} = \vec{0}$. That is, for each $n \in \{0, 1, \dots, M\}$, the equation $2x_n + x_{2^n} + x_{2^{n+1}} + \cdots + x_{2^{n+1}-1} = y_n$ holds.

Let $r \geq M$ and assume that we have chosen $\langle x_i \rangle_{i=0}^{2^{r+1}-1}$ and $\langle y_i \rangle_{i=0}^r$ in C such that for each $n \in \{0, 1, \dots, r\}$, the equation $2x_n + x_{2^n} + x_{2^{n+1}} + \cdots + x_{2^{n+1}-1} = y_n$ holds. For $2^{r+1} \leq t \leq 2^{r+1} + m - 3$, let $x_t = x_{r+1}$. Let $k = 2^{r+1} - m + 2$. Then $k \geq K$ and

$$2x_{r+1} + x_{2^{r+1}} + x_{2^{r+1}+1} + \cdots + x_{2^{r+1}+m-3} = m \cdot x_{r+1} \in m \cdot \mathbb{Z}$$

so by Lemma 1.4 we may pick $y_{r+1} \in C$ and pick $x_t \in C$ for $2^{r+1} + m - 2 \leq t \leq 2^{r+2} - 1$ such that $2x_{r+1} + x_{2^{r+1}} + x_{2^{r+1}+1} + \cdots + x_{2^{r+2}-1} = y_{r+1}$. \square

We shall see in Corollary 3.8 that in fact one may choose \vec{x} and \vec{y} in C^ω such that $(A \ -I) \begin{pmatrix} \vec{x} \\ \vec{y} \end{pmatrix} = \vec{0}$ and all entries of $\begin{pmatrix} \vec{x} \\ \vec{y} \end{pmatrix}$ are distinct.

The matrix of Theorem 2.1 has unbounded row sums. This motivates the following. Let us say that an $\omega \times \omega$ matrix A with entries from \mathbb{Z} has *bounded row sums* if $\{\sum_{j<\omega} |a_{i,j}| : i < \omega\}$ is bounded.

We now see that if A is a matrix which is partition regular and has bounded row sums then in fact some non-empty set of columns must sum to zero. In fact, this is pretty much a direct copy of Rado's original proof in the finite case.

Theorem 2.2. *Let A be an $\omega \times \omega$ matrix with entries from \mathbb{Z} having bounded row sums. If A is KPR/ \mathbb{N} and \vec{c}_i denotes column i of A , then there is some $J \subseteq \omega$ such that $\sum_{i \in J} \vec{c}_i = \vec{0}$.*

Proof. Pick a prime q such that $\sum_{j<\omega} |a_{i,j}| < q$ for each $i < \omega$. Given $x \in \mathbb{N}$, pick $b(x) \in \{1, 2, \dots, q-1\}$, $l(x) \in \omega$ and $a(x) \in \omega$ such that $x = b(x)q^{l(x)} + a(x)q^{l(x)+1}$. (Thus $b(x)$ is the rightmost nonzero digit in the base q expansion of x and $l(x)$ is the position of that digit.) For $b \in \{1, 2, \dots, q-1\}$, let $C_b = \{x \in \mathbb{N} : b(x) = b\}$. Pick $b \in \{1, 2, \dots, q-1\}$ and $\vec{x} \in C_b^\omega$ such that $A\vec{x} = \vec{0}$. Let $d = \min\{l(x_i) : i < \omega\}$ and let $J = \{i < \omega : l(x_i) = d\}$. We shall show that $\sum_{j \in J} \vec{c}_j = \vec{0}$.

For $j \in J$, let $e_j = a(x_j)$ so that $x_j = bq^d + e_jq^{d+1}$. For $j \in \omega \setminus J$, $l(x_j) > d$ so pick $e_j \in \mathbb{N}$ such that $x_j = e_jq^{d+1}$. Suppose that $\sum_{j \in J} \vec{c}_j \neq \vec{0}$ and pick some $i < \omega$ such that $\sum_{j \in J} a_{i,j} \neq 0$. Then $0 = \sum_{j \in J} bq^d a_{i,j} + \sum_{j < \omega} e_jq^{d+1} a_{i,j}$ so q divides $\sum_{j \in J} ba_{i,j}$ so q divides $\sum_{j \in J} a_{i,j}$ while $q > \sum_{j \in J} |a_{i,j}|$, a contradiction. \square

Remark 2.3. A matrix A which satisfies the weaker condition that $\{|\sum_{j<\omega} a_{i,j}| : i < \omega\}$ is bounded could be partition regular and have no non-empty set of columns summing to zero. If

$$A = \begin{pmatrix} -2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & \cdots \\ 0 & -3 & 1 & 1 & 0 & 0 & 0 & 0 & \cdots \\ 0 & 0 & -4 & 0 & 1 & 1 & 1 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

then $(A \ -I)$ is partition regular; the proof is essentially the same as that of Theorem 2.1. Clearly, there is no non-empty set of columns of $(A \ -I)$ whose sum is zero.

Very frustratingly, we have been unable to extend Theorem 2.2 to the following.

Conjecture 2.4. *Let A be an $\omega \times \omega$ matrix with entries from \mathbb{Z} , and assume that A is partition regular. If A has bounded row sums, then A has the columns property.*

If this conjecture is false, might at least a weaker statement be true, that A must satisfy the natural 'transfinite' version of the columns property?

Definition 2.5. Let $u, v \in \mathbb{N} \cup \{\omega\}$ and let A be a $u \times v$ matrix with entries from \mathbb{Q} . Denote the columns of A by $\langle \vec{c}_i \rangle_{i < v}$. The matrix A satisfies the *transfinite columns property* if and only if there exist a countable ordinal μ and a partition $\langle I_\sigma \rangle_{\sigma < \mu}$ of v such that

$$(1) \sum_{i \in I_0} \vec{c}_i = \vec{0} \text{ and}$$

- (2) for each $t \in \mu \setminus \{0\}$, $\sum_{i \in I_t} \vec{c}_i$ is a linear combination with coefficients from \mathbb{Q} of $\{\vec{c}_i : i \in \bigcup_{j < t} I_j\}$.

We remark that, for any given countable ordinal μ , it is a straightforward exercise to write down a matrix that has the transfinite columns property with the partition being indexed by μ , but without any partition having a smaller indexing ordinal.

Our weaker version of Conjecture 2.4 would then be:

Conjecture 2.6. *Let A be an $\omega \times \omega$ matrix with entries from \mathbb{Z} and assume that A is partition regular. If A has bounded row sums, then A has the transfinite columns property.*

We saw in the Introduction that the system of equations $x_n - x_{n+1} = y_n$ satisfies the columns property but is not partition regular over \mathbb{N} . However, as shown at the end of Section 3 of [6], it is partition regular over \mathbb{R} — meaning that whenever $\mathbb{R} \setminus \{0\}$ is finitely coloured there is a monochromatic solution to that system of equations. (It was also shown there that it is not partition regular over \mathbb{Q} .) This leaves open the possibility that perhaps a sufficiently well behaved system of equations which satisfies the columns property must at least be partition regular over \mathbb{R} . To end this section, we show that this seems not to be the case. The system $x_n + 2x_{n+1} = y_n$ is about as well behaved as one can wish. The sum of absolute values of entries of each row of the coefficient matrix is 4 and all columns sum to 1, -1 , or 3. The matrix equation is

$$\begin{pmatrix} 1 & -1 & 2 & 0 & 0 & 0 & 0 & 0 & \cdots \\ 0 & 0 & 1 & -1 & 2 & 0 & 0 & 0 & \cdots \\ 0 & 0 & 0 & 0 & 1 & -1 & 2 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \\ x_1 \\ y_1 \\ \vdots \end{pmatrix} = \vec{0}.$$

The matrix of coefficients satisfies the columns property with $I_i = \{2i, 2i + 1\}$ for each $i < \omega$.

Theorem 2.7. *The system of equations $x_n + 2x_{n+1} = y_n$ is not partition regular over \mathbb{R} .*

Proof. We show first that the system is not partition regular over \mathbb{Q} , for which it suffices to show that it is not partition regular over $\mathbb{Q}^+ = \{x \in \mathbb{Q} : x > 0\}$. We begin by defining $\tau : \mathbb{Q}^+ \rightarrow \{0, 1, 2\}$ by $\tau(x) = i$ if and only if $\lfloor \log_2(x) \rfloor \equiv i \pmod{3}$. Note that if $\tau(x) = \tau(y)$ and $y \geq 2x$, then $y > 4x$. (To see this, let $k = \lfloor \log_2(x) \rfloor$ and $m = \lfloor \log_2(y) \rfloor$. Then $2^{m+1} > y \geq 2x \geq 2^{k+1}$ so $m > k$ and therefore $m \geq k + 3$. Then $y \geq 2^m \geq 2^{k+3} > 4x$.)

As is relatively well known, and at any rate easy to verify, every rational $x \in (0, 1)$ has a unique expansion of the form

$$x = \sum_{t=2}^{m(x)} \frac{a(x, t)}{t!}$$

where each $a(x, t) \in \{0, 1, \dots, t-1\}$ and $a(x, m(x)) > 0$. If $t > m(x)$, let $a(x, t) = 0$. For each $t \geq 2$, choose $\nu_t : \{1, 2, \dots, t-1\} \rightarrow \{0, 1, 2\}$ such that, if $i, j \in \{1, 2, \dots, t-1\}$ and $j \equiv 2i \pmod{t}$, then $\nu_t(i) \neq \nu_t(j)$. (To see that such a colouring exists, let G be the digraph on vertex set $\{1, 2, \dots, t-1\}$ with an edge

from i to j whenever $j \equiv 2i \pmod{t}$. Then every edge of G is in at most one cycle, so G can be 3-coloured.)

Now define a finite colouring φ of \mathbb{Q}^+ so that for $x, y \in \mathbb{Q}^+$, $\varphi(x) = \varphi(y)$ if and only if

- (1) $\tau(x) = \tau(y)$;
- (2) $x < 1$ if and only if $y < 1$; and
- (3) if $x < 1$ and $y < 1$ then
 - (a) $m(x) \equiv m(y) \pmod{3}$ and
 - (b) if $m(x) = m(y)$, then $\nu_{m(x)}(a(x, m(x))) = \nu_{m(y)}(a(y, m(y)))$.

Now suppose we have a system $x_n + 2x_{n+1} = y_n$ for $n < \omega$ such that $\{x_n : n < \omega\} \cup \{y_n : n < \omega\}$ is monochromatic with respect to φ . Given n , we have that $y_n > 2x_{n+1}$ so, since $\tau(y_n) = \tau(x_{n+1})$, we have $y_n > 4x_{n+1}$. But then $x_n = y_n - 2x_{n+1} > 2x_{n+1}$. Therefore the sequence $\langle x_n \rangle_{n=0}^\infty$ is eventually less than 1 and therefore $\{x_n : n < \omega\} \cup \{y_n : n < \omega\} \subseteq (0, 1)$. Given any k , $\{x \in (0, 1) : m(x) \leq k\}$ is finite and therefore, there is some n such that $m(x_{n+1}) > m(x_n)$ and consequently $m(x_{n+1}) \geq m(x_n) + 3$. Let $m = m(x_{n+1})$, let $(d, c, b) = (a(x_{n+1}, m-2), a(x_{n+1}, m-1), a(x_{n+1}, m))$, and let $(d', c', b') = (a(y_n, m-2), a(y_n, m-1), a(y_n, m))$. Now $b' \equiv 2b \pmod{m}$ so, since $\varphi(y_n) = \varphi(x_{n+1})$ we must have that $b' = 0$ and therefore $m(y_n) < m(x_{n+1}) = m$. Thus $m(y_n) \leq m - 3$ and so $d' = c' = 0$. Since $0 \equiv 2b \pmod{m}$ we must have that $m = 2b$. Since there is a carry out of position m when $2x_{n+1}$ is computed, $0 = c' \equiv 2c + 1 \pmod{m-1}$ and so there is a carry out of position $m-1$. But then $0 = d' \equiv 2d + 1 \pmod{m-2}$, which is impossible since $m-2$ is even.

Thus we have established that our system is not partition regular over \mathbb{Q}^+ . Extend the colouring φ to a colouring φ' of $\mathbb{Q} \setminus \{0\}$ with respect to which there is no monochromatic system (e.g. by reflecting to $\{x \in \mathbb{Q} : x < 0\}$ using a new set of colours).

Now pick a Hamel basis B for \mathbb{R} over \mathbb{Q} and well-order B . For $x \in \mathbb{R} \setminus \{0\}$ let $\text{supp}(x)$ be the finite non-empty subset of B such that $x = \sum_{b \in \text{supp}(x)} \alpha(x, b)b$ and each $\alpha(x, b) \in \mathbb{Q} \setminus \{0\}$ and let $b(x) = \max \text{supp}(x)$.

Now let $\psi : \mathbb{Q} \setminus \{0\} \rightarrow \{0, 1\}$ such that for all $x \in \mathbb{Q} \setminus \{0\}$, $\psi(2x) \neq \psi(x)$. (For example, colour by the parity of the number of factors of 2 in the numerator or denominator of x .) Now define a finite colouring γ of $\mathbb{R} \setminus \{0\}$ so that $\gamma(x) = \gamma(y)$ if and only if

- (1) $\alpha(x, b(x)) > 0$ if and only if $\alpha(y, b(y)) > 0$;
- (2) $\psi(\alpha(x, b(x))) = \psi(\alpha(y, b(y)))$; and
- (3) $\varphi'(\alpha(x, b(x))) = \varphi'(\alpha(y, b(y)))$.

Now suppose we have a system $x_n + 2x_{n+1} = y_n$ in \mathbb{R} for $n < \omega$ such that $\{x_n : n < \omega\} \cup \{y_n : n < \omega\}$ is monochromatic with respect to γ . If for any n we have $b(x_{n+1}) > b(x_n)$, then we get $b(y_n) = b(x_{n+1})$ and $\alpha(y_n, b(y_n)) = 2\alpha(x_{n+1}, b(x_{n+1}))$, contradicting requirement (2). Therefore, for each n , $b(x_{n+1}) \leq b(x_n)$. Since there are no infinite strictly decreasing sequences in B , there are some $b \in B$ and $k < \omega$ such that for all $n \geq k$, $b(x_n) = b$. Note that by requirement (1), for all such n , $b(y_n) = b$. Now let for each $n < \omega$, $x'_n = \alpha(x_{k+n}, b)$ and $y'_n = \alpha(y_{k+n}, b)$. Then $\{x'_n : n < \omega\} \cup \{y'_n : n < \omega\}$ is monochromatic with respect to φ' and for each n , $x_n + 2x_{n+1} = y_n$, a contradiction. \square

3. PROOF OF LEMMA 1.4

In this section we will provide a proof of a generalisation of Lemma 1.4. We also prove a strengthening of Theorem 2.1. These results utilise the algebraic structure of the Stone–Čech compactification βS of a discrete semigroup S . We shall provide a brief introduction to this structure now. For proofs of the assertions made here, see the first four chapters of [10]. We shall be concerned here almost exclusively with commutative semigroups, so we will denote the operation of S by $+$. We mention here that the reader who is not interested in the results for general groups or semigroups can skip Lemma 3.1 and its proof and rejoin the text at Lemma 3.3, taking G to be \mathbb{Z} and d to be the usual density for subsets of \mathbb{Z} .

We take the points of βS to be the ultrafilters on S , the principal ultrafilters being identified with the points of S , whereby we pretend that $S \subseteq \beta S$. Given a subset A of S , $\overline{A} = \{p \in \beta S : A \in p\}$ is the closure of A in βS and $\{\overline{A} : A \subseteq S\}$ is a basis for the open sets of βS as well as a basis for the closed sets of βS .

The operation on S extends to an operation on βS , also denoted by $+$, with the property that for each $p \in \beta S$, the function $\rho_p : \beta S \rightarrow \beta S$ is continuous and for each $x \in S$, the function $\lambda_x : \beta S \rightarrow \beta S$ is continuous where, for $q \in \beta S$, $\rho_p(q) = q + p$ and $\lambda_x(q) = x + q$. (The reader should be cautioned that $(\beta S, +)$ is almost certain to not be commutative; the centre of $(\beta\mathbb{N}, +)$ is \mathbb{N} .) Given $A \subseteq S$ and $p, q \in \beta S$, $A \in p + q$ if and only if $\{x \in S : -x + A \in q\} \in p$, where $-x + A = \{y \in S : x + y \in A\}$.

Thus $(\beta S, +)$ is a compact Hausdorff right topological semigroup. As with any such object, there exist idempotents in βS . Also, βS has a smallest two sided ideal, $K(\beta S)$, which is the union of all of the minimal right ideals of βS as well as the union of all of the minimal left ideals of βS . The intersection of any minimal right ideal with any minimal left ideal is a group and any two such groups are isomorphic. In particular, there are idempotents in $K(\beta S)$. A subset C of S is said to be *central* if and only if it is a member of some idempotent in $K(\beta S)$. (Such idempotents are said to be *minimal*.) Since whenever the union of finitely many sets is a member of an ultrafilter, one of the sets must itself be a member of the ultrafilter, one has immediately that whenever S is finitely coloured, at least one colour class must be central. A subset A of S is said to be *central** if and only if it has non-empty intersection with any central set, equivalently if and only if it is a member of every minimal idempotent.

Given a set X , let $\mathcal{P}_f(X)$ be the set of finite non-empty subsets of X . A subset A of S is *syndetic* if and only if there is some $L \in \mathcal{P}_f(S)$ such that $S = \bigcup_{t \in L} (-t + A)$. A subset A of S is *piecewise syndetic* if and only if

$$(\exists G \in \mathcal{P}_f(S)) (\forall F \in \mathcal{P}_f(S)) (\exists x \in S) (F + x \subseteq \bigcup_{t \in G} (-t + A)).$$

It is then a fact that A is piecewise syndetic if and only if $\overline{A} \cap K(\beta S) \neq \emptyset$. In particular, every central set is piecewise syndetic.

Lemma 3.1. *Let $(S, +)$ be a commutative cancellative semigroup. There is a function $d : \mathcal{P}(S) \rightarrow [0, 1]$ with the following properties.*

- (1) $d(S) = 1$.
- (2) For each piecewise syndetic subset A of S , $d(A) > 0$.
- (3) For $B, C \subseteq S$, $d(B \cup C) \leq d(B) + d(C)$.
- (4) For $a \in S$ and $B \subseteq S$, $d(B) = d(a + B) = d(-a + B)$.
- (5) If $B \subseteq S$, $n \in \mathbb{N}$, and $x_1, x_2, \dots, x_n \in S$ such that $(x_i + B) \cap (x_j + B) = \emptyset$ if $i \neq j$, then $d(\bigcup_{i=1}^n (x_i + B)) = nd(B)$.

Proof. There are at least two ways to see this. One way is to use the following very general result, whose proof can be found in [3].

Lemma 3.2. *Let S be a left amenable semigroup and let A be a piecewise syndetic subset of S . Then a probability measure ν can be defined on βS with the following properties:*

- (a) $\nu(s^{-1}B) = \nu(B)$ for every Borel subset B of βS and every $s \in S$ and
- (b) $\nu(\overline{A}) > 0$.

If, in addition S is cancellative, then:

- (c) $\nu(sB) = \nu(B)$ for every Borel subset B of βS and every $s \in S$.

Using this lemma, one then defines $d(B) = \sup\{\nu(\overline{B}) : \nu \text{ is a probability measure on } \beta S \text{ satisfying (a) and (c) of Lemma 3.2}\}$.

An alternative, mostly elementary, method is to use the fact proved by Argabright and Wilde in [1, Theorem 4] that any commutative semigroup satisfies the *Strong Følner Condition*:

$$(\forall H \in \mathcal{P}_f(S)) (\forall \epsilon > 0) (\exists K \in \mathcal{P}_f(S)) (\forall s \in H) (|K \Delta (s + K)| < \epsilon \cdot |K|)$$

(For an entirely elementary proof of this fact see [8, Section 7].)

If S satisfies the Strong Følner Condition and $A \subseteq S$, then the *Følner density* of A is

$$d(A) = \sup \left\{ \alpha : (\forall H \in \mathcal{P}_f(S)) (\forall \epsilon > 0) (\exists K \in \mathcal{P}_f(S)) (|A \cap K| \geq \alpha \cdot |K|) \text{ and } (\forall s \in H) (|K \Delta (s + K)| < \epsilon \cdot |K|) \right\}.$$

Conclusions (1) and (3) are routine to verify for Følner density and conclusion (4) is established in [7, Theorem 4.17]. Define

$$\mathbf{D} = \{p \in \beta S : (\forall A \in p)(d(A) > 0)\}.$$

To verify (2), let A be a piecewise syndetic subset of S . By [9, Theorems 2.12, 2.14, and 5.9], \mathbf{D} is a two-sided ideal of βS . (Alternatively, it is a routine elementary argument to establish directly that \mathbf{D} is a two-sided ideal of βS .) Since \mathbf{D} is a two-sided ideal of βS , $K(\beta S) \subseteq \mathbf{D}$. Since A is piecewise syndetic, $\overline{A} \cap K(\beta S) \neq \emptyset$ and therefore $\overline{A} \cap \mathbf{D} \neq \emptyset$. Therefore $d(A) > 0$.

Finally, it is routine to establish that conclusion (5) holds. To emphasize how routine it is, we present the full proof now. By conclusions (3) and (4), $d(\bigcup_{i=1}^n (x_i + B)) \leq \sum_{i=1}^n d(x_i + B) = nd(B)$. Let $a = d(B)$. We complete the proof by showing that if $\alpha < a$, then $d(\bigcup_{i=1}^n (x_i + B)) \geq n\alpha$.

Let $\alpha < a$ be given and let $\gamma = (a - \alpha)/2$. To see that $d(\bigcup_{i=1}^n (x_i + B)) \geq n\alpha$, let $H \in \mathcal{P}_f(S)$ and $\epsilon > 0$ be given. Let $H' = H \cup \{x_1, x_2, \dots, x_n\}$ and let $\epsilon' = \min\{\epsilon, \gamma\}$. Since $d(B) > \alpha + \gamma$, pick $K \in \mathcal{P}_f(S)$ such that $|B \cap K| \geq (\alpha + \gamma)|K|$ and $(\forall s \in H') (|K \Delta (s + K)| < \epsilon'|K|)$. Then $(\forall s \in H) (|K \Delta (s + K)| < \epsilon|K|)$ so it suffices to show that $|(\bigcup_{i=1}^n (x_i + B)) \cap K| \geq n\alpha|K|$. By the disjointness assumption, $|(\bigcup_{i=1}^n (x_i + B)) \cap K| = \sum_{i=1}^n |(x_i + B) \cap K|$ so it suffices to let $i \in \{1, 2, \dots, n\}$ and show that $|(x_i + B) \cap K| \geq \alpha|K|$.

So let $i \in \{1, 2, \dots, n\}$ be given. Since $x_i \in H'$, we have that $|(x_i + K) \setminus K| < \epsilon'|K|$. Therefore

$$\begin{aligned} |(x_i + B) \cap K| &\geq |(x_i + B) \cap (x_i + K)| - |(x_i + K) \setminus K| \\ &> |B \cap K| - \epsilon'|K| \\ &\geq (\alpha + \gamma)|K| - \gamma|K| \\ &= \alpha|K|. \end{aligned} \quad \square$$

For the rest of this section we will assume we have a density function d as guaranteed by Lemma 3.1.

The next few lemmas are based on very similar results from [2]. Recall that, given a commutative semigroup $(S, +)$, $A \subseteq S$ and $k \in \mathbb{N}$, $kA = A + A + \dots + A$ (k times) and $k \cdot A = \{k \cdot s : s \in A\}$.

Lemma 3.3. *Let $(G, +)$ be a commutative group and let A be a piecewise syndetic subset of G . Then $A - A$ is syndetic in G .*

Proof. Pick $H \in \mathcal{P}_f(G)$ such that

$$(\forall F \in \mathcal{P}_f(G))(\exists y \in G)(F + y \subseteq \bigcup_{t \in H} (-t + A)).$$

Let $L = H - H$. We claim that $G \subseteq \bigcup_{s \in L} (-s + (A - A))$. To see this, let $x \in G$. Pick $y \in G$ such that $\{0, x\} + y \subseteq \bigcup_{t \in H} (-t + A)$. Pick t_1 and t_2 in H such that $t_1 + x + y \in A$ and $t_2 + 0 + y \in A$. Then $x + (t_1 - t_2) \in A - A$. \square

Lemma 3.4. *Let $(G, +)$ be a commutative group and let $S \subseteq G$ such that $0 \in S$, $S = -S$, and $d(S) > 0$. Then there is a subgroup E of G such that if $l \geq 2/d(S)$, then $lS = E$.*

Proof. We will show that there is some $j \leq 1/d(S)$ such that $(2j+1)S = (2j)S$, and so $(k+1)S = kS$ for $k \geq 2/d(S)$. Once we have shown this, let $k = \lceil 2/d(S) \rceil$ and let $E = kS$. We have that $E + E = kS + kS = (2k)S = E$. Since S is symmetric, we also have that $E = -E$, so E is closed under addition and the taking of inverses, hence is a subgroup of G as required.

Suppose instead that for each $j \leq 1/d(S)$, $(2j)S \subsetneq (2j+1)S$. We claim that, for each such j , $(2j+1)S$ contains $j+1$ disjoint translates of S . This is a contradiction for $j = \lfloor 1/d(S) \rfloor$.

The claim is true for $j = 0$. For $j > 0$, choose $x \in (2j+1)S \setminus (2j)S$. Then $x = s_1 + \dots + s_{2j+1}$ with $s_i \in S$ for each i . We have

$$\begin{aligned} S + x - s_1 &= S + s_2 + \dots + s_{2j+1} \subseteq (2j+1)S \\ \text{and } (2j-1)S - s_1 &\subseteq (2j)S \subseteq (2j+1)S. \end{aligned}$$

Since $(2j-1)S - s_1$ contains j disjoint translates of S , it suffices to show that $S + x - s_1$ and $(2j-1)S - s_1$ are disjoint. But if they intersect then $t_0 + x - s_1 = t_1 + \dots + t_{2j-1} - s_1$ for some $t_i \in S$, from which it follows that $x = t_1 + \dots + t_{2j-1} - t_0 \in (2j)S$, contradicting the choice of x . \square

Lemma 3.5. *Let $(G, +)$ be a commutative group and let $S \subseteq G$ such that $0 \in S$ and $d(S) > 0$. Then there exists $Y \subseteq G$ such that for $k \geq 2/d(S)$, we have $S - kS = Y$.*

Proof. We suppose to the contrary that $S - (2j)S \subsetneq S - (2j+1)S$ for all $j \leq 1/d(S)$ and show that $S - (2j+1)S$ contains $j+1$ disjoint translates of S for each such j , which is impossible for $j = \lfloor 1/d(S) \rfloor$.

The claim is true for $j = 0$. For $j > 0$, choose $x \in (S - (2j + 1)S) \setminus (S - (2j)S)$. Then $x = s_0 - s_1 - \dots - s_{2j+1}$ with $s_i \in S$ for each i . We have

$$S + x - s_0 \subseteq S - (2j + 1)S, \text{ and} \\ S - (2j - 1)S - s_0 \subseteq S - (2j)S \subseteq S - (2j + 1)S,$$

hence it suffices to show that $S + x - s_0$ and $S - (2j - 1)S - s_0$ are disjoint. But if they intersect then $t_0 + x - s_0 = t_1 - t_2 - \dots - t_{2j} - s_0$ for some $t_i \in S$, whence $x = t_1 - t_2 - \dots - t_{2j} - t_0 \in S - (2j)S$, contradicting the choice of x . \square

Lemma 3.6. *Let $(G, +)$ be a commutative group and let $A \subseteq G$ such that $d(A) > 0$. Assume that E is a subgroup of G such that if $l \geq 2/d(A - A)$, then $l(A - A) = E$. If $k \geq 2/d(A)$, then $(A - kA) = (A - kA) + E$.*

Proof. Let $k = \lceil 2/d(A) \rceil$ and let $X = A - kA$. For any $a \in A$, $0 \in A - a$ and $d(A - a) = d(A)$, so by Lemma 3.5, if $Y = (A - a) - k(A - a)$ then also $Y = (A - a) - (k + 1)(A - a)$ and so $X = X - (A - a)$. Letting a range over the whole of A gives $X = X + (A - A)$. Then for all $l \in \mathbb{N}$, $X = X + l(A - A)$, hence $X = X + E$ as required. \square

The following is our promised generalisation of Lemma 1.4.

Lemma 3.7. *Let $(G, +)$ be a commutative group and assume that $n \cdot G$ is a central* set for each $n \in \mathbb{N}$. Let A be a central subset of G . Then $d(A) > 0$ and there exists $m \in \mathbb{N}$ such that if $k \geq 2/d(A)$, then $m \cdot G \subseteq A - kA$.*

Proof. Since A is piecewise syndetic, by Lemma 3.1 $d(A) > 0$. Let $S = A - A$. Pick by Lemma 3.4 a subgroup E of G such that if $l \geq 2/d(S)$, then $lS = E$. By Lemma 3.6 we have that if $k \geq 2/d(A)$, then $(A - kA) = (A - kA) + E$.

Now $A - A$ is syndetic by Lemma 3.3. Let $l = \lceil 2/d(A - A) \rceil$. Then $A - A \subseteq l(A - A) = E$, so E is syndetic. Pick $n \in \mathbb{N}$ and $x_1, x_2, \dots, x_n \in G$ such that $G = \bigcup_{i=1}^n (-x_i + E)$. For $i \in \{1, 2, \dots, n\}$ pick $a_i < b_i$ in \mathbb{N} and $j \in \{1, 2, \dots, n\}$ such that $a_i \cdot x_i \in (-x_j + E)$ and $b_i \cdot x_i \in (-x_j + E)$. Then $(b_i - a_i) \cdot x_i \in E - E = E$. Let $m = \prod_{i=1}^n (b_i - a_i)$. Then for each $i \in \{1, 2, \dots, n\}$, $m \cdot x_i \in E$.

We claim that $m \cdot G \subseteq E$. So let $x \in G$ and pick $i \in \{1, 2, \dots, n\}$ such that $x \in (-x_i + E)$. Then $x_i + x \in E$ so $m \cdot x_i + m \cdot x \in E$. Since $m \cdot x_i \in E$, we have that $m \cdot x \in E$.

Finally, let $k \geq 2/d(A)$. Then $(A - kA) = (A - kA) + E$. Since $m \cdot G$ is central*, $A \cap m \cdot G \neq \emptyset$. If $x \in A \cap m \cdot G$, then $x - k \cdot x \in (A - kA) \cap E$ so $E = (x - k \cdot x) + E \subseteq A - kA$. Thus $m \cdot G \subseteq A - kA$. \square

Proof of Lemma 1.4. Let C be a central subset of \mathbb{N} . By [10, Exercise 4.3.8] $K(\beta\mathbb{Z}) = K(\beta\mathbb{N}) \cup -K(\beta\mathbb{N})$. Therefore C is a central subset of \mathbb{Z} . Given $n \in \mathbb{N}$, by [10, Lemma 6.6], $n \cdot \mathbb{N}$ is a member of every idempotent in $\beta\mathbb{N}$, so in particular is a member of every idempotent in $K(\beta\mathbb{N})$. Again using the fact that $K(\beta\mathbb{Z}) = K(\beta\mathbb{N}) \cup -K(\beta\mathbb{N})$, we then have that $n \cdot \mathbb{Z}$ is central* in \mathbb{Z} . Therefore Lemma 3.7 applies. \square

We are now ready for our strengthening of Theorem 2.1.

Corollary 3.8. *Let A be the matrix of Theorem 2.1. For any central subset C of \mathbb{N} , there exist \vec{x} and \vec{y} in C^ω such that $(A \quad -I) \begin{pmatrix} \vec{x} \\ \vec{y} \end{pmatrix} = \vec{0}$ and all entries of $\begin{pmatrix} \vec{x} \\ \vec{y} \end{pmatrix}$ are distinct.*

Proof. Let C be a central set and pick an idempotent $p \in K(\beta\mathbb{N})$ such that $C \in p$. Let $C^* = \{x \in C : -x + C \in p\}$. By [10, Lemma 4.14], if $x \in C^*$, then $-x + C^* \in p$. By Lemma 3.7, pick $m \in \mathbb{N}$ such that for all $k \geq 2/d(C^*)$, $m \cdot \mathbb{Z} \subseteq C^* - kC^*$. Pick $M \in \mathbb{N}$ such that $2^{M+1} \geq m - 2 + \lceil 2/d(C^*) \rceil$.

Let P be the $(M+1) \times 2^{M+1}$ matrix consisting of rows $0, 1, \dots, M$ and columns $0, 1, \dots, 2^{M+1} - 1$ of A and let I_{M+1} be the $(M+1) \times (M+1)$ identity matrix. If the columns of $\begin{pmatrix} I_{M+1} \\ P \end{pmatrix}$ are reversed, then the first nonzero entry in each column is 1 so by [10, Corollary 15.6], $\begin{pmatrix} I_{M+1} \\ P \end{pmatrix}$ is image partition regular over \mathbb{N} and thus by [10, Theorem 15.24(m)] we may choose $\vec{x} = \langle x_i \rangle_{i=0}^{2^{M+1}-1}$ and $\vec{y} = \langle y_i \rangle_{i=0}^M$ in C^* such that all entries of $\begin{pmatrix} \vec{x} \\ \vec{y} \end{pmatrix}$ are distinct.

Now let $r \geq M$ and assume that we have chosen $\langle x_i \rangle_{i=0}^{2^{r+1}-1}$ and $\langle y_i \rangle_{i=0}^r$ in C^* such that for each $n \in \{0, 1, \dots, r\}$, $y_n = 2x_n + \sum_{j=2^n}^{2^{n+1}-1} x_n$ and all of the x_i and y_j are distinct. For $2^{r+1} \leq t \leq 2^{r+2} - 1$, let $z_t = x_{r+1}$. Let $k = 2^{r+1} - m + 2$. Then $k \geq \lceil 2/d(C^*) \rceil$ and $2x_{r+1} + \sum_{t=2^{r+1}}^{2^{r+1}+m-3} z_t = m \cdot x_{r+1} \in m \cdot \mathbb{Z}$ so by Lemma 3.7 we may pick $v \in C^*$ and pick $z_t \in C^*$ for $2^{r+1} + m - 2 \leq t \leq 2^{r+2} - 1$ such that $v = 2x_{r+1} + \sum_{t=2^{r+1}}^{2^{r+2}-1} z_t$. Let

$$D = (-v + C^*) \cap \bigcap_{t=2^{r+1}}^{2^{r+2}-1} (-z_t + C^*).$$

Then $D \in p$ which is an idempotent so by [10, Theorem 5.8] pick a sequence $\langle w_n \rangle_{n=1}^\infty$ in \mathbb{N} such that $FS(\langle w_n \rangle_{n=1}^\infty) = \{\sum_{t \in F} x_t : F \in \mathcal{P}_f(\mathbb{N})\} \subseteq D$. We may assume that the sequence $\langle w_n \rangle_{n=1}^\infty$ is increasing. Choose $n(2^{r+1}) \in \mathbb{N}$ such that $w_{n(2^{r+1})} > \max(\{x_i : i \in \{0, 1, \dots, 2^{r+1} - 1\}\} \cup \{y_i : i \in \{0, 1, \dots, r\}\})$. Given $j \in \{2^{r+1}, 2^{r+1} + 1, \dots, 2^{r+2} - 2\}$, having chosen $n(j)$, pick $n(j+1) > n(j)$ such that $w_{n(j+1)} > z_j + w_{n(j)}$.

For $2^{r+1} \leq j \leq 2^{r+2} - 1$, let $x_j = z_j + w_{n(j)}$ and let $y_{r+1} = 2x_{r+1} + \sum_{j=2^{r+1}}^{2^{r+2}-1} x_j$. Then each $x_j \in C^*$ and $y_{r+1} = v + \sum_{j=2^{r+1}}^{2^{r+2}-1} w_{n(j)} \in C^*$. Finally, $\max(\{x_i : i \in \{0, 1, \dots, 2^{r+1} - 1\}\} \cup \{y_i : i \in \{0, 1, \dots, r\}\}) < x_{2^{r+1}} < x_{2^{r+1}+1} < \dots < x_{2^{r+2}-1} < y_{r+1}$. \square

Remark 3.9. We remark that Corollary 3.8 holds more generally. Let G be a commutative group with the property that, for every $n \in \mathbb{N}$, $n \cdot G$ is a central subset of G . Then Corollary 3.8 holds with \mathbb{N} replaced by G . The proof is essentially the same.

REFERENCES

1. L. Argabright and C. Wilde, *Semigroups satisfying a strong Følner condition*, Proc. Amer. Math. Soc. **17** (1967), 587–591.
2. B. Barber, N. Hindman, and I. Leader, *Partition regularity in the rationals*, J. Comb. Theory (Series A) **120** (2013), 1590–1599.
3. H. Dales, A. Lau, and D. Strauss, *Banach algebras on semigroups and on their compactifications*, Mem. Amer. Math. Soc. **205** (2010), 165 pp.
4. N. Hindman, *Finite sums from sequences within cells of a partition of \mathbb{N}* , J. Comb. Theory (Series A) **17** (1974), 1–11.

5. N. Hindman, I. Leader, and D. Strauss, *Image partition regular matrices – solutions in central sets*, Trans. Amer. Math. Soc. **355** (2003), 1213–1235.
6. N. Hindman, I. Leader, and D. Strauss, *Open problems in partition regularity*, Combin. Probab. Comput. **12** (2003), 571–583.
7. N. Hindman and D. Strauss, *Density in arbitrary semigroups*, Semigroup Forum **73** (2006), 273–300.
8. N. Hindman and D. Strauss, *Sets satisfying the Central Sets Theorem*, Semigroup Forum **79** (2009), 480–506.
9. N. Hindman and D. Strauss, *Density and invariant means in left cancellative left amenable semigroups*, Topology Appl. **156** (2009), 2614–2628.
10. N. Hindman and D. Strauss, *Algebra in the Stone–Čech compactification: theory and applications, 2nd edition*, Walter de Gruyter & Co., Berlin, 2012.
11. K. Milliken, *Ramsey’s Theorem with sums or unions*, J. Comb. Theory (Series A) **18** (1975), 276–290.
12. R. Rado, *Studien zur Kombinatorik*, Math. Z. **36** (1933), 424–470.
13. I. Schur, *Über die Kongruenz $x^m + y^m = z^m \pmod{p}$* , Jahresbericht der Deutschen Math.-Verein. **25** (1916), 114–117.
14. A. Taylor, *A canonical partition relation for finite subsets of ω* , J. Comb. Theory (Series A) **21** (1976), 137–146.
15. B. van der Waerden, *Beweis einer Baudetschen Vermutung*, Nieuw Arch. Wiskunde **19** (1927), 212–216.

SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, EDGBASTON, BIRMINGHAM, B15 2TT, UK.

E-mail address: `b.a.barber@bham.ac.uk`

DEPARTMENT OF MATHEMATICS, HOWARD UNIVERSITY, WASHINGTON, DC 20059, USA.

E-mail address: `nhindman@aol.com`

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, CENTRE FOR MATHEMATICAL SCIENCES, WILBERFORCE ROAD, CAMBRIDGE, CB3 0WB, UK.

E-mail address: `i.leader@dpms.cam.ac.uk`

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9J2, UK.

E-mail address: `d.strauss@hull.ac.uk`